# CENTURY-SCALE (1886 – 2018) BLUFF EROSION: BLOCK ISLAND, RHODE ISLAND, USA



Bryan A. Oakley
Environmental Earth Science Department
Eastern Connecticut State University

**2022 NEGSA ANNUAL MEETING: T13. Current Research in Coastal and Marine Processes** 




# Block Island, Rhode Island



#### Block Island, Rhode Island

- 25 km<sup>2</sup> Island located 15 km south of mainland RI
- Bluffs vary in height from <2.5 m to >
   50 m in relief
- Two segments of moraine connected by barrier spits (complicated stratigraphy)
- Microtidal (<1 m) wave-dominated shoreline</li>



#### Stratigraphy of a tectonic end moraine Block Island (Mohegan Bluffs)


MO = Meltout Till

OW = Outwash (tectonized w/ older till (LD) incorporated

LDL = Older Lakefloor deposits (varves) (tectonized)

LD = Older till – in place??

Older "Lower Drift" - (Early Wisconsinan? Illinoian?)



## Stratigraphy of a tectonic end moraine Block Island (Side view)






# **Block Island: Post Sandy (June 2013)**





# **Block Island: Pre and Post Sandy LiDAR**





# Block Island: June 2015








# Block Island: Slump, circa 2000






J. Boothroyd photo; in Kelley (20XX)

## **Block Island: Spring 2019 Slump**






# **Block Island: Spring 2019 Slump**

#### **Block Island: Shoreline Change Maps**



-13.03 m



# Data and Methods: Uncertainty


$$E_{sp} = V(E_g^2 + E_d^2 + E_t^2 + E_l^2)$$

Measurement Uncertainties (meters)	1886 T-Sheet	2018 LiDAR
Georeferencing (E <sub>g</sub> )	2.8	N/A
Digitizing (E <sub>d</sub> )	1	1
T-sheet survey (E <sub>t</sub> )	10	N/A
Lidar position uncertainty (E <sub>I</sub> )	N/A	1
Total position uncertainty (E <sub>sp</sub> )	10.4	1.4
Annualized retreat rate uncertainty (m yr <sup>-1</sup> )	0.08	

#### **Data and Methods**


Bluff crest and shoreline digitized in both 1886 and 2018 LiDAR

#### 1886 T-Sheet




#### **Data and Methods**

- Transects Cast at 25 m spacing (n = 630 transects with a bluff)
- Net Shoreline Movement measured 1886 – 2018 for both bluff crest and shoreline





#### Results

- 1.2% of Transects showed 'accretion' (n = 8) along gullies
- 86% of transects (n = 545) exceeded the positional uncertainty
- Mean retreat 28 m;
   maximum retreat 98 m




# Results: Clay Head (Northeast shoreline)







### **Results: Southern Shoreline**



Results:
Southeast
&
Southwest







# **Results: Obligatory Data Table**

Location	# of Transects	Ave Bluff Height (m)	Ave Bluff NSM (m)	Ave Bluff EPR (m yr <sup>-1</sup> )	Ave Shoreline NSM (m)	Ave Shoreline EPR (m yr <sup>-1</sup> )
Grove Point to  Mansion Beach	101	20.8	-31.9	-0.24	-38.2	-0.29
Scotch Beach	25	8.6	-26.3	-0.20	-44.9	-0.34
Old Harbor to Southeast Point	104	14.9	-20.4	-0.15	-21.7	-0.17
Southeast Point to Schooner Point	225	30.0	-30.4	-0.23	-37.8	-0.29
Schooner Point to Grace Cove	167	11.4	-28.5	-0.22	-25.3	-0.19
Overall	622	20.1	-28.3	-0.21	-32.1	-0.25


## Results: Obligatory Graph – Shoreline vs. Bluff Retreat




# Results: Bluff Height v. Bluff Retreat



#### **Results: Transect Orientation v. Retreat**





#### **Conclusions and Future Work**

- Bluff Retreat for Block Island varied between 'no change' and >98 m
- Some slight variation in average retreat for different areas of the island (generally



#### **Conclusions and Future Work**

- Comparing the bluff retreat to shoreline retreat showed.....
- No clear trend with Bluff Height and Retreat



#### **Conclusions and Future Work**

- Finish this part of the project examining other aspects affecting retreat
- Continue examining processes using LiDAR and SfM

